

# 1° EXERCISE


- Two long straight , parallel wires, 12.5 cm apart, carry current of equal magnitude “I”.

They repel each other with a force per unit length of  $7.6 \times 10^{-9}$  N/m.

Are the currents parallel or anti parallel. Find “I”.

# S\_1° The current are antiparallel

Forza magnetica tra due fili percorsi da corrente



Magnetic field at wire 2 from current in wire 1:

$$B = \frac{\mu_0 I_1}{2\pi r}$$

Force on a length  $\Delta L$  of wire 2:

$$F = I_2 \Delta L B$$

Force per unit length in terms of the currents:

$$\frac{F}{\Delta L} = \frac{\mu_0 I_1 I_2}{2\pi r}$$

## 2° EXERCISE

- A wire of length 22cm floats 1.5mm above a long straight wire.
- The same current flows in both wire.
- If the mass of the 22cm wire is 8g, what is the current?

$$S_2^\circ \quad F = P$$

- $$F = k \frac{i_1 i_2 l}{d} = m^* g$$

# 3° EXERCISE

- A 32cm straight, horizontal wire of mass 80g is connected to a source of emf by flexible leads.
- A magnetic field of 2T is perpendicular to the wire.
- Find the current necessary to let the wire float.

- $F = I * L * B * \text{sen}(\theta) = m * g$

- $I = ( m * g ) / ( L * B * \text{sen}(\theta) ) =$

$$= ( 0.080 * 9.81 ) / ( 0.32 * 2 * \text{sen}(90^\circ) )$$

# 4° EXERCISE

- An electron moving with velocity “ $v$ ” to the right enters a region of uniform magnetic field that point out of the page.
- After the electron enter this region, will it be deflected upward or downward?

- 

UPWARD